



Volume contents

Volume 46 (1994)

Issue contents	1	ment of wing hair polarity ends prior to the start of hair morphogenesis		
M.J. Sonnenfeld and J.R. Jacobs			101	
Mesectodermal cell fate analysis in Drosophila midline mutants	3	J.F. de Celis, A. García-Bellido Roles of the <i>Notch</i> gene in <i>Drosophila</i> wing morph genesis		
A. Macías, S. Pelaz and G. Morata		Be	109	
Genetic factors controlling the expression of the abdominal-A gene of Drosophila within its domain D.I. Strutt and R.A.H. White	15	M. Lardelli, J. Dahlstrand, U. Lendahl The novel Notch homologue mouse Notch 3 lac specific epidermal growth factor-repeats and expressed in proliferating neuroepithelium		
Characterisation of T48, a target of homeotic gene regulation in <i>Drosophila</i> embryogenesis	27	J. Jorre, G.B.L.J. van der Lans, P.H. Lanser, J.M.A. Vervaart, D. Zivkovic, J.E. Speksnijder, W. Kruijer		
D. Pan, S.A. Valentine and A.J. Courey The bipartite D. melanogaster twist promoter is reorganized in D. virilis	41	Effects of retinoic acid on the expression of retinoic acid receptors during zebrafish embryogenesis	137	
C. Walsh, A. Glaser, R. Fundele, A. Ferguson-Smith, S. Barton, M.A. Surani and R. Ohlsson		Issue contents	151	
The non-viability of uniparental mouse conceptuses correlates with the loss of the products of imprinted genes	55	E. Sánchez-Herrero, I. Guerrero, J. Sampedro, A. González-Reyes Developmental consequences of unrestricted expres-		
ML. Dirksen, M.I. Morasso, T.D. Sargent and M. Jamrich		sion of the abd-A gene of Drosophila	153	
Differential expression of a Distal-less homeobox gene Xdll-2 in ectodermal cell lineages	63	M. Rothe, E.A. Wimmer, M.J. Pankratz, M. González-Gaitán, H. Jäckle Identical transacting factor requirement for knirps and knirps-related gene expression in the anterior but		
Issue contents	71	not in the posterior region of the <i>Drosophila</i> embryo	169	
C. Wülbeck, C. Fromental-Ramain, J.A. Campos-Ortega The HLH domain of a zebrafish HE12 homologue can partially substitute for functions of the HLH		M. González-Gaitán, M.P. Capdevila, A. García-Bellido Cell proliferation patterns in the wing imaginal disc of <i>Drosophila</i>	183	
domain of <i>Drosophila</i> DAUGHTERLESS	73	A W. Carlon		
J.C. Ruiz, E.J. Robertson The expression of the receptor-protein tyrosine kinase gene, eck, is highly restricted during early		A.W. Stoker Isoforms of a novel cell adhesion molecule-like protein tyrosine phosphatase are implicated in neural development	201	
mouse development	87			
P.N. Adler, J. Charlton, K.H. Jones, J. Liu The cold-sensitive period for frizzled in the develop-		R.S. Winning, T.D. Sargent Pagliaccio, a member of the Eph family of receptor tyrosine kinase genes, has localized expression in a		

subset of neural crest and neural tissues in Xenopus laevis embryos	219	Author index	249
D. Kostic, M.R. Capecchi	219	Subject index	251
Targeted disruptions of the murine <i>Hoxa-4</i> and <i>Hoxa-6</i> genes result in homeotic transformations of components of the vertebral column	231	Volume contents	253

